

Robotic surgery for gastric cancer: expert opinion from a western perspective

Amilcare Parisi

St. Mary's Hospital, Department of Digestive Surgery,
Terni, Italy.

Gastric surgery is one of the most relevant fields of development for minimally invasive technologies. Laparoscopy is now widespread, and several studies have demonstrated its feasibility and safety even in some advanced oncological procedures. Robotic surgery has several intrinsic advantages that theoretically can improve an extensive lymphadenectomy or the reconstruction phase.

Much remains to be addressed in this field and further studies are necessary to offer the patient the best possible approach based on his characteristics and the stage of his disease. The present report offers an overview on robotics and its role in gastric surgery.

Authors

*Amilcare Parisi*¹

¹ St. Mary's Hospital, Department of Digestive Surgery, Terni, Italy.

To Cite

Parisi A. Robotic surgery for gastric cancer: expert opinion from a western perspective. *J Gastric Surg* 2019; 1(1): 16-18

Publication history

Received: November 6, 2019

Revised: November 28, 2019

Accepted: December 2, 2019

Article in press: December 10, 2019

Published online: December 16, 2019

Correspondence to

Dr. Amilcare Parisi

Department of Digestive Surgery,

St Mary's Hospital, Terni 05100, Italy

Telephon: +393387312313

Background

Robotic technology has spread throughout the world in the last decades and Italy has been one of the first countries introducing robotic systems in Europe.

In recent years the number of procedures has dramatically increased.

Over 86 robotic systems are currently active, and the number of procedures is now over 13,000 operations per year. Therefore, nowadays, Italy together with France is the first country in Europe for the total number of robotic surgical procedures performed per year.

Robotic technology has revolutionized our approach to minimally invasive surgery.

In this context, many fields of interest are currently under development: new devices and platforms, imaging integrated technology, teaching opportunities, new techniques.

The Xi robot represents the 4th generation of the Da Vinci systems and the evolution of the previous Si version. Moreover, new devices have been produced with instruments even more advanced. For example, the endowristed robotic stapler that allows the surgeon to sectioning by a correct angle of vision from the console. The vessel sealer uses advanced radiofrequency energy and could be particularly useful for liver parenchyma transection.

Then, the single site surgery could open the possibility for new applications even for gastric surgery.

Today the Xi system offers the possibility of combining images from different sources. For example, it's possible to perform an intraoperative ultrasound or follow a gastroscopy, from the robotic console, together with the 3D view of the operative field. These developments are improving and changing the way we perform oncological surgical procedures and new concepts of surgery are now emerging: the navigation surgery and the targeted surgery[1].

Robotic technology represents also a huge opportunity for education. Young surgeons can perform different surgical steps of complex procedures through mentoring. Robotic simulators are also available.

So, training in robotic surgery can start at the simulator and then go through the execution of different surgical steps from the easier to the challenging ones, and finally approach the entire operation.

Robotic gastric surgery

Certainly, oncologic gastric surgery is one of the areas of greatest interest and development in the field of minimally invasive and robotic surgery[2-4].

Many centers have published their own experiences[5] and comparative studies, giving authors of systematic reviews and meta-analysis the opportunity to try to define the role of robotic surgery by comparing it with traditional laparoscopy or open surgery. However, the current level of evidence is far from being able to consider these procedures in common surgical practice. Current guidelines describe laparoscopy as a possible alternative to open surgery for early gastric cancer, while robotic surgery is recognized to have some advantages, but this approach is not mentioned in gastric cancer guidelines[6].

The main discussed limitations are the proper execution

of an extended lymphadenectomy, when required, and a safe approach for reconstruction. The latter is still the object of controversy and most surgeons are concerned about the possibility to

perform a totally intracorporeal procedure.

Face the reconstruction with new approaches: the Parisi's Technique

The technique we have developed for reconstruction after total gastrectomy is called "double loop method"[7-9] in which a first loop of bowel is measured and shifted up antecolic to directly perform the esophago-jejunal anastomosis. The jejunal loop is secured to the posterior wall of the esophagus with seromuscular interrupted stitches (second posterior layer of the anastomosis). The row of metallic stitches at the esophagus margin is removed and a 3 cm incision is made at the jejunum side. A double continuous suturing is performed with PDS 3/0. The anastomosis is completed with a second anterior layer performed with interrupted stitches.

The second loop (alimentary limb) is measured up to 40 cm starting from the esophago-jejunostomy and is fixed to the biliary limb close to the previous anastomosis on its left side.

A jejuno-jejunal anastomosis is now performed as usual. Finally, the two anastomoses are interrupted by firing the linear stapler.

Preliminary results of this approach were previously reported[7, 9].The double loop method increases the feasibility of a full hand-sewn intracorporeal reconstruction.

Robotic Gastrectomy in Italy: IMIGASTRIC data

We have looked to the data collected in the IMIGASTRIC database[10-13] and we extracted those patients who underwent robotic surgery in Italian Institutions to show their characteristics and outcomes.

Overall data of 222 patients were found. The average age was 68yo, BMI around 25, 50% of patients had comorbidities. At the pathology examination 56% of cases were Advanced Gastric Cancer, most with undifferentiated histology. Majority of patients, 74%, underwent a distal gastrectomy.

Among the intraoperative outcomes, mean EBL was 137ml, a low rate of intraoperative complications occurred, there were no major intraoperative complications or death.

Overall, during the postoperative course, patients underwent mobilization on the 1 postoperative day, the liquid diet was administered at the 3 POD and a soft solid diet on the 4 POD. Intravenous analgesics were used for an average of 3 days. Mean hospital stay resulted 8 days. 16% of patients experienced medical or surgical related postoperative complications. The majority were low- grade complications (CD I+II), while only 5 patients (2%) had major complications requiring a reoperation. 4% of patients were readmitted after discharge. No death occurred. Among surgical complications, 6 patients had leakages (leakage rate 2,7%) but only one underwent reoperation.

Conclusion

Despite the potential benefits of robotic surgery, there is still a lot of research to be done in this field and the scientific community is asking what strategies need to be adopted in future studies to develop and standardize minimally invasive surgery.

That's why, 3 years ago, we have established a multi- institutional cooperation with a project aimed to collect eastern and western data in a large registry, called IMIGASTRIC. A network is today needed to bring together different experiences and institutions with the aim of improving medical

and surgical care in gastric cancer patients.

Acknowledgements

Not applicable.

Contributors

AP was involved in designing the study, analyzing the literature, references searching and in writing the final manuscript.

Funding

None.

Competing interests

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval

Not applicable

Provenance and peer review

Not commissioned; externally peer reviewed.

Open access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non- Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work noncommercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

References

1. Desiderio J, Trastulli S, Gemini A, Di Nardo D, Palazzini G, Parisi A, et al. Fluorescence image-guided lymphadenectomy using indocyanine green and near infrared technology in robotic gastrectomy. *Chin J Cancer Res.* 2018;30:568-70..
2. Tu RH, Lin JX, Desiderio J, Li P, Xie JW, Wang JB, et al. Does Intra- Abdominal Infection after Curative Gastrectomy Affect Patients' Long- Term Prognosis? A Multi-Center Study Based on a Large Sample Size. *Surg Infect (Larchmt).* 2019;20:271-7..
3. Lin JX, Lin JP, Desiderio J, Xie JW, Gemini A, Wang JB, et al. Difference in the short-term outcomes of laparoscopic gastrectomy for gastric carcinoma between the east and west: a

- retrospective study from the IMIGASTRIC trial. *J Cancer*. 2019;10:4106-13..
4. Huang ZN, Desiderio J, Chen QY, Zheng CH, Li P, Xie JW, et al. Indications for adjuvant chemotherapy in patients with AJCC stage IIa T3N0M0 and T1N2M0 gastric cancer-an east and west multicenter study. *BMC Gastroenterol*. 2019;19:205..
 5. Desiderio J, Stewart CL, Sun V, Melstrom L, Warner S, Lee B, et al. Enhanced Recovery after Surgery for Gastric Cancer Patients Improves Clinical Outcomes at a US Cancer Center. *J Gastric Cancer*. 2018;18:230-41..
 6. Marano A, Choi YY, Hyung WJ, Kim YM, Kim J, Noh SH. Robotic versus Laparoscopic versus Open Gastrectomy: A Meta-Analysis. *Journal of gastric cancer*. 2013;13:136-48..
 7. Parisi A, Ricci F, Gemini A, Trastulli S, Cirocchi R, Palazzini G, et al. New totally intracorporeal reconstructive approach after robotic total gastrectomy: Technical details and short-term outcomes. *World J Gastroenterol*. 2017;23:4293-302..
 8. Parisi A, Trastulli S, Ricci F, Cirocchi R, Pironi D, Santoro A, et al. Robotic double-loop reconstruction method following total gastrectomy. *Endoscopy*. 2016;48 Suppl 1 UCTN:E55-6..
 9. Parisi A, Ricci F, Trastulli S, Cirocchi R, Gemini A, Grassi V, et al. Robotic Total Gastrectomy With Intracorporeal Robot-Sewn Anastomosis: A Novel Approach Adopting the Double-Loop Reconstruction Method. *Medicine (Baltimore)*. 2015;94:e1922..
 10. Parisi A, Reim D, Borghi F, Nguyen NT, Qi F, Coratti A, et al. Minimally invasive surgery for gastric cancer: A comparison between robotic, laparoscopic and open surgery. *World J Gastroenterol*. 2017;23:2376-84..
 11. Parisi A, Nguyen NT, Reim D, Zhang S, Jiang ZW, Brower ST, et al. Current status of minimally invasive surgery for gastric cancer: A literature review to highlight studies limits. *Int J Surg*. 2015;17:34-40..
 12. Parisi A, Desiderio J. Establishing a multi-institutional registry to compare the outcomes of robotic, laparoscopic, and open surgery for gastric cancer. *Surgery*. 2015;157:830-1..
 13. Desiderio J, Jiang ZW, Nguyen NT, Zhang S, Reim D, Alimoglu O, et al. Robotic, laparoscopic and open surgery for gastric cancer compared on surgical, clinical and oncological outcomes: a multi-institutional chart review. A study protocol of the International study group on Minimally Invasive surgery for GASTRIC Cancer-IMIGASTRIC.. *BMJ Open*. 2015;5:e008198..